



Sample and Carrier




In each case several carrier glasses are analyzed with always new application of material onto the carrier, not all images came from one and the same mRNA application. Several times (10x) mRNA was applied of several carrier for all the images as shown.








Reflected light MicType1

MRNA sample



MRNA sample









4-8 μm diameter/ thickness (not exactly measurable)






Very fine hair = 20 - 40 μm common 50 - 80 μm

=> At least 2 to 5 times thinner than very fine hair






Reflected light MicTyp2







Transmitted light MicTyp2

Blood Reference




Blood Reference



Optical comparison examples Campra

Tratamiento: Exfoliación de escamas de rGO mediante 30s sonicación suave con pulsos de 5s mediante sonda



A small, irregularly shaped, light-colored object, possibly a piece of paper or a small animal, is centered on a dark, textured background. The object has a mottled appearance with shades of light brown and beige. The background is dark and grainy, with some faint, lighter-colored streaks or fibers visible. The overall image is somewhat blurry and has a low resolution.


This image shows a blank page with a light beige or off-white background. There are several small, dark specks scattered across the surface, which appear to be dust or scanning artifacts. The overall texture is slightly grainy, typical of a scanned document page.

Please contact a sales office, sales representative or distributor for more information. http://www.oxfordjournals.org/academic/oxfordjournals/oxfordjournals.org	
French: <i>Par</i> O: <i>FR00A</i>	Public Company Issued 1578-0707(200601)13:1;1-2  1578-0707(200601)13:1;1-2
	F: <i>en</i> M: <i>JM</i>

PATRON DE OXIDO DE GRAFENO REDUCIDO



MUESTRA
RD1



DETECCIÓN DE OXIDO DE GRAFENO
EN SUSPENSIÓN ACUOSA (COMIRNATY™ (RD1))


ESTUDIO OBSERVACIONAL EN MICROSCOPIA ÓPTICA Y ELECTRÓNICA

Informe provisional (I)
28 de Junio de 2021






Prof. Dr. Pablo Campra Madrid
Doctor en Ciencias Químicas y Licenciado en Ciencias Biológicas
ESCUELA SUPERIOR DE INGENIERÍA
UNIVERSIDAD DE ALMERÍA, ESPAÑA



MUESTRA
RD1







Optical comparison examples Campra



MRNA analyzed sample











PATRON DE OXIDO DE GRAFENO REDUCIDO 10X

Tratamiento: Exfoliación de escamas de rGO mediante 30s sonicación suave con pulsos de 5s mediante sonda



DETECCION DE OXIDO DE GRAFENO
EN SUSPENSION ACUOSA (CONFIRMADO: rGO)

ESTUDIO OBSERVACIONAL EN MICROSCOPIA OPTICA Y ELECTRONICA

Informe provisional (I)

28 de junio de 2023



Prof. Dr. Pablo Campa Madrid


Doctor en Ciencias Químicas y Licenciado en Ciencias Biológicas

ESPECIALIDAD DE QUÍMICA

UNIVERSIDAD DE ZARAGOZA, ESPAÑA



MUESTRA
RD1



MUESTRA
RD1 60x

MUESTRA
RD1
(campo
oscuro)



Optical comparison examples Campra

MRNA analyzed sample


Sample with cover glass

Edge cover glass

Glass carrier

Edge cover glass
leaked sample

100 μ m



10 μ m

PATRÓN DE ÓXIDO DE GRAFENO REDUCIDO 10X

Tratamiento: Exfoliación de escamas de rGO mediante 30s sonicación suave con pulsos de 5s mediante sonda



DETECCIÓN DE ÓXIDO DE GRAFENO
EN SUSPENSIÓN ACUOSA (CONFIRMACIÓN: IPI)

ESTUDIO OBSERVACIONAL EN MICROSCOPIA ÓPTICA Y ELECTRÓNICA


Informe provisional (I)

28 de junio de 2023




Prof. Dr. Pablo Campa Madrid

Doctor en Ciencias Químicas y Licenciado en Ciencias Biológicas
ESPECIALIDAD DE FÍSICA
UNIVERSIDAD DE JAÉN, ESPAÑA




MUESTRA
RD1




MUESTRA
RD1 60x


MUESTRA
RD1
(campo
oscuro)




Optical comparison examples Campra



20 µm



20 µm



20 µm

PATRON DE OXIDO DE GRAFENO REDUCIDO 10X

Tratamiento: Exfoliación de escamas de rGO mediante 30s sonicación suave con pulsos de 5s mediante sonda



DETECCIÓN DE ÓXIDO DE GRAFENO
EN SUSPENSIÓN ACUOSA (CONFIRMACIÓN: I)

ESTUDIO OBSERVACIONAL EN MICROSCOPIA ÓPTICA Y ELECTRÓNICA

Informe provisional (I)

28 de junio de 2023




Prof. Dr. Pablo Campa Madrid

Departamento de Química y Física de Materiales, Centro de Investigación en Nanotecnología,
Escuela Superior de Ingeniería
UNIVERSIDAD DE ZARAGOZA, ESPAÑA




MUESTRA
RD1






MUESTRA
RD1 60x



MUESTRA
RD1
(campo
oscuro)




Optical comparison examples Campra




MRNA analyzed sample (only structure with different color appearance)




Dried samples from previous analysis (dried 14 days with glass cover then cover removed), small residual amount of liquid present)




Dried samples from previous analysis (dried 14 days with glass cover then cover removed), small residual amount of liquid present)




Dried samples from previous analysis (dried 14 days with glass cover then cover removed), small residual amount of liquid present)





Dried samples from previous analysis (dried 14 days with glass cover then cover removed), small residual amount of liquid present)





Dried samples from previous analysis (dried 14 days with glass cover then cover removed), small residual amount of liquid present)



Dried samples from previous analysis (dried 14 days with glass cover then cover removed), small residual amount of liquid present)




Dried samples from previous analysis (dried 14 days with glass cover then cover removed), small residual amount of liquid present)





dried samples from
previous analysis (dried
14 days with glass cover
then cover removed)







Transition glass liquid and glass only as a reference of a clean carrier





Darkfield Mic3





















From two
images put
together
the
complete
structure





























Examination with mixing blood and transmission
field microscopy Mic2

Analysis Procedure:

- Glasscarrier and glass cover cleaned with isopropanol and cleanroom wipes, additionally cleaned with compressed air
- Prick from finger and remove from first blood leakage
- Blood drop on glass slide one part added with ModeRNA vaccine
- ModeRNA quantity corresponds to approx. 1/20 of the administered dose and is mixed with the blood in a ratio of approx. 1:1 (eye measure)
- Reference blood and that with blood+ModeRNA cover applied as fast as possible
- Standing time in air until covered maximum 20s (standing time is slightly longer than normal as shortest possible standing time is desired)
- Bottle of ModeRNA has been stored at room temperature.
- Images were taken with a transmitting light microscope, magnification 5-100x as needed

Something similar was shown with Dr. Flemming:


<https://www.flemingmethod.com/the-pfizer-vaccine-blood>

Left blood not covered, middle only blood as reference, right blood with ModeRNA


It is already obvious that the flow with ModeRNA is less good, although ModeRNA alone flows very well in earlier analysis alone.

The left site was then also treated with ModeRNA and covered.




(the poorer flow behavior may also be due to the slightly longer process, see better performed in dark field part)



The dot marks approx. where the recordings are from






Blood Reference






The dot marks approx. where the recordings are from

Blood + Moderna




This can be seen in part also with NACL




Destroyed platelets, this can be seen partly also in normal blood towards the edge

A second attempt was made to see if it is reproducible with the same procedure.







Here, too, the sample with ModeRNA no longer flowed properly, even optically one can see a kind of clotting with the normal camera.





Blood

Blood reference
different positions








Where blood accumulates and cannot flow further (end of glass cap), it looks like the picture on the left below.



Blood




Blood reference
several places




At the point where the blood accumulates and cannot flow further (end of glass cap) it looks like in the pictures below. However, it should not confuse with the effect of blood with MRNA, where the blood did not flowed or has flowed little from the center.



Blood






Also with normal blood it can look like shown, however, this was so evident only at the edge



Blood + Moderna





Several positions




Blood + Moderna





Several positions




Blood + Moderna





Several positions




Blood + Moderna






Several positions




Blood + Moderna





several positions,
more outside of the
blood center




Blood + Moderna








several positions,
more outside of the
blood center



NACL "NAPREEP" for reference




NACL „Serophy“ as Reference




Comparison Blood+NACL and Blood+ModeRNA

Person had little water intake (i.e. no water for last 4h)







Person had 0.7l water intake 30min before



Especially when you had drunk little water MRNA flows almost not, with water a little better but generally worse than with NACL




Blood+ModeRNA

Blood+ MRNA




Blood+ModeRNA




Blood+ MRNA






Again this large structure from the MRNA



Blood+ NACL







Blood+ NACL







Examination with mixing blood
Dark field Mic3

Blood with granulocytes, still easily observable, with a lot of movement, few prickly apple shapes, blood carrier stored directly next to Blood+MRNA carrier. The blood has already strongly degraded, but still assessable.



The same sample is always viewed at different time





Here, too, the blood is still easily assessable, even granulocytes are still present. Healthy blood can "live" on the carrier and be assessed for up to 10 days if stored correctly.



The same sample is always viewed at different times

Here you can see how the mixed blood has already clotted, the blood is no longer assessable. Various particles from the vaccine are also visible here.

Blood + MRNA (Mixture ca. 1:5)




The same sample was observed at different times

Blood + MRNA (Mixture ca. 1:5)

15.12.2021 9:10

Blood disintegrates, many large gaps without fillite, blood static, without movement, agglomerations.
No granulocytes visible.



The same sample is always viewed at different times

Summary transmitted light analysis

- The analyzed glass carrier has been shown to be clean and shows no structures or impurities which could explain the seen contamination/structures
- Structures seen in the Swiss sample that cannot be defined what they are based on the ingredients known
- These structures correspond to the pictures and videos of other analyzed samples from different countries.
- There are sharp-edged structures (rectangles and the like mostly in the range of 0.5 - 75 μ m)
- There are elongated round structures which have a diameter of about 6-8 μ m and a length up to 1mm.
- There are wrinkle-like structures and according to the pictures and videos of other analyses of other analyzed samples from different countries
- The structures themselves are very thin (assumption is 100-500nm) and mostly difficult to get into focus in the liquid.
- There are structures that look like small rocks which are according to the pictures and videos of similar analyses of Covid vaxx samples from different countries
- There are structures like small rings
- The difference in structures would either have to be based on a broad contamination base which is not expected in such productions or it is intended in there. It would have been rather expected to find uniform structures like small spheres or repetitive structures but not as in the variety and variation seen

Summary darkfield analysis

- With the darkfield microscopy the images of the transmission light microscopy can be shown from a different point of view
- In each analysis with new sample carrier and new application of MRNA material these long round structures can be seen (mostly 1-2x per carrier/application of MRNA which is quite frequent)
- With darkfield nothing could be seen that was not possible to be seen with transmitted light microscopy, the with transmitted light quality of the images was even better

Summary blood analysis transmission light microscopy

- With a 1:1 mixture with the clots seem to have formed especially if one had drunk little water prior. Can this be a factor of more serious side effects, people who drink rather little or had not drunk well before vaccination?
- The time covering the mixed probe lasted a few seconds longer and may have an influence, but this would not be expected in this extreme form
- Also with NACL trial has shown similar effects as with MRNA but less strong, here is also referred to the video of Flemming which explains this but for laymen the difference seems not fully clear
- These effects have also been shown with Flemming:
(<https://www.flemingmethod.com/the-pfizer-vaccine-blood>)Diese Effekte
- Here the question arises what it means when you have 1:1 mixture and to what extent this can occur in the body in a similar way causing harm

Summary blood analysis darkfield

- Blood with granulocytes was well observable, with a lot of movement, few datura forms
- Such samples can be kept and further analyzed for several days if stored well (Both samples reference blood and the mixed sample were stored together in the same place next to each other)
- After half a day the blood sample blood was still easily assessable, even granulocytes were still present
- Blood mixed with MRNA already clotted from the beginning although the blood standing time was kept as short in comparison to the the transmitted light analysis, the blood was immediately no longer assessable.
- Various particles from the vaccine are also visible here as well as this long structure
- Blood mixed with MRNA disintegrates strongly after half a day, many large gaps without fillite, the blood is static and without movement with agglomerations, no granulocytes visible anymore

General Conclusion

- The structures were compared with the Campra graphene oxide and reduced graphene oxide images and correspond to these very well whether it is this or not cannot be said by microscopy alone
- Additional structure analysis is required for a definite statement
- Visually it is a good indication based on the reports available worldwide, but not sufficient proof, also the Campra Mirco Raman analysis needs to be redone once more to prove its correctness
- Support for the interpretation of the images or support for further analyses (SEM/TEM, EDX, micro Raman or further suggestions) is requested
- For the blood analyses direct feedback is requested

